

Polynomial Representation of E_6 and Its Combinatorial and PDE Implications¹

Xiaoping Xu

Institute of Mathematics, Academy of Mathematics & System Sciences

Chinese Academy of Sciences, Beijing 100190, P.R. China ²

Abstract

In this paper, we use partial differential equations to find the decomposition of the polynomial algebra over the basic irreducible module of E_6 into a sum of irreducible submodules. It turns out that the cubic polynomial invariant corresponding to the Dicksons' invariant trilinear form is the unique fundamental invariant. Moreover, we obtain a combinatorial identity saying that the dimensions of certain irreducible modules of E_6 are correlated by the binomial coefficients of twenty-six. Furthermore, we find all the polynomial solutions for the invariant differential operator corresponding to the Dickson trilinear form in terms of the irreducible submodules.

1 Introduction

The E_6 Lie algebra and group are popular mathematical objects with broad applications. Dickson [13] (1901) first realized that there exists an E_6 -invariant trilinear form on its 27-dimensional basic irreducible module. The 78-dimensional simple Lie algebra of type E_6 can be realized by all the derivations and multiplication operators with trace zero on the 27-dimensional exceptional simple Jordan algebra (e.g., cf. [1], [30]). Aschbacher [5] used the Dickson form to study the subgroup structure of the E_6 group. Bion-Nadal [8] proved that the E_6 Coxeter graph can be realized as a principal graph of subfactor of the hyperfinite Π_1 factor. Brylinski and Kostant [9] obtained a generalized Capelli identity on the minimal representation of E_6 . Binegar and Zierau [7] found a singular representation of E_6 . Ginzburg [16] proved that the twisted partial L -function on the 27-dimensional

¹2000 Mathematical Subject Classification. Primary 17B10, 17B25; Secondary 17B01.

²Research supported by China NSF 10871193

representation of $GE_6(\mathbb{C})$ is entire except the points 0 and 1. Iltyakov [21] showed that the field of invariant rational functions of E_6 on the direct sum of finite copies of the basic module and its dual is purely transcendental. Suzuki and Wakui [29] studied the Turaev-Viro-Oceanu invariant of 3-manifolds derived from the E_6 -subfactor. Moreover, Cerchiai and Scotti [11] investigated the mapping geometry of the E_6 group. Furthermore, the (A_2, G_2) duality in E_6 was obtained by Rubenthaler [28].

Okamoto and Marshak [27] constructed a grand unification preon model with E_6 metacolor. The E_6 Lie algebra was used in [18] to explain the degeneracies encountered in the genetic code as the result of a sequence of symmetry breakings that have occurred during its evolution. Wang [31] identified Geoner's model with the twisted LG model and E_6 singlets. Morrison, Pieruschka and Wybourne [26] constructed the E_6 interacting boson model. Berglund, Candelas et al. [6] studied instanton contributions to the masses and couplings of E_6 singlets. Haba and Matsuoka [17] found large lepton flavor mixing in the E_6 -type unification models. Ghezelbash, Shafiekhani and Abolbasani [15] derived explicitly a set of Picard-Fuchs equations of $N = 2$ supersymmetric E_6 Yang-Mills theory. Anderson and Blažek [2-4] found certain Clebsch-Gordan coefficients in connection with the E_6 unification model building. Fernández-Núñez, Garcia-Fuertes and Perelomov [14] used the quantum Calogero-Sutherland model corresponding to the root system of E_6 to calculate Clebsch-Gordan series for this algebra. Howl and King [20] proposed a minimal E_6 supersymmetric standard model which allows Planck scale unification, provides a solution to the μ problem and predicts a new Z' . Das and Laperashvili [12] studied Preon model related to family replicated E_6 unification.

The motivation of this work is to understand the functional impact of the E_6 Lie algebra (group). We use partial differential equations to decompose the polynomial algebra over the basic irreducible module of E_6 into a sum of irreducible submodules. Consequently, the cubic polynomial invariant corresponding to the Dickson's invariant trilinear form is the unique fundamental invariant (any other invariant is a polynomial in it). Moreover, we obtain a combinatorial identity, which says that the dimensions of certain irreducible modules of E_6 are correlated by the binomial coefficients of twenty-six. Furthermore, we find all the polynomial solutions for the invariant differential operator corresponding to the Dickson trilinear form in terms of the irreducible submodules. Below we give a more detailed introduction to our results.

It has been known for many years that the representation theory of Lie algebra is closely related to combinatorial identities. Macdonald [25] generalized the Weyl denominator

identities for finite root systems to those for infinite affine root systems, which are now known as the Macdonald's identities. We present here a consequence of the Macdonald's identities taken from Kostant's work [22]. Let $\mathcal{G}$ be a finite-dimensional simple Lie algebra over the field $\mathbb{C}$ of complex numbers. Denote by Λ^+ the set of dominant weights of $\mathcal{G}$ and by $V(\lambda)$ the finite-dimensional irreducible $\mathcal{G}$ -module with highest weight λ . It is known that the Casimir operator takes a constant $c(\lambda)$ on $V(\lambda)$. Macdonald's Theorem implies that there exists a map $\chi : \Lambda^+ \rightarrow \{-1, 0, 1\}$ such that the following identity holds:

$$\left(\prod_{n=1}^{\infty} (1 - q^n)\right)^{\dim \mathcal{G}} = \sum_{\lambda \in \Lambda^+} \chi(\lambda) (\dim V(\lambda)) q^{c(\lambda)}. \quad (1.1)$$

Kostant [23] connected the above identity to the abelian subalgebras of $\mathcal{G}$.

Denote by $E_{r,s}$ the square matrix with 1 as its (r, s) -entry and 0 as the others. The orthogonal Lie algebra

$$o(n, \mathbb{C}) = \sum_{1 \leq r < s \leq n} \mathbb{C}(E_{r,s} - E_{s,r}). \quad (1.2)$$

It acts on the polynomial algebra $\mathcal{A} = \mathbb{C}[x_1, \dots, x_n]$ by

$$(E_{r,s} - E_{s,r})|_{\mathcal{A}} = x_r \partial_{x_s} - x_s \partial_{x_r}. \quad (1.3)$$

Denote by $\mathcal{A}_k$ the subspace of homogeneous polynomials in $\mathcal{A}$ with degree k . When $n \geq 3$, it is well known that the subspace of harmonic polynomials

$$\mathcal{H}_k = \{f \in \mathcal{A}_k \mid (\partial_{x_1}^2 + \dots + \partial_{x_n}^2)(f) = 0\} \quad (1.4)$$

forms an irreducible $o(n, \mathbb{C})$ -module and

$$\mathcal{A}_k = \mathcal{H}_k \oplus (x_1^2 + x_2^2 + \dots + x_n^2) \mathcal{A}_{k-2}. \quad (1.5)$$

Recall the special linear Lie algebra

$$sl(n, \mathbb{C}) = \sum_{r \neq s} \mathbb{C} E_{r,s} + \sum_{r=1}^{n-1} \mathbb{C}(E_{r,r} - E_{r+1,r+1}). \quad (1.6)$$

Let $\mathcal{A}$ be the polynomial algebra in $x_1, \dots, x_n, y_1, \dots, y_n$. Define a representation of $sl(n, \mathbb{C})$ on $\mathcal{A}$ via

$$E_{r,s}|_{\mathcal{A}} = x_r \partial_{x_s} - y_s \partial_{y_r}. \quad (1.7)$$

Denote by $\mathbb{N}$ the additive semigroup of nonnegative integers. Define

$$x^\alpha = x_1^{\alpha_1} x_2^{\alpha_2} \cdots x_n^{\alpha_n} \quad \text{for } \alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n. \quad (1.8)$$

Set

$$\mathcal{A}_{\ell_1, \ell_2} = \sum_{\alpha, \beta \in \mathbb{N}^n; |\alpha|=\ell_1, |\beta|=\ell_2} \mathbb{R} x^\alpha y^\beta \quad \text{for } \ell_1, \ell_2 \in \mathbb{N} \quad (1.9)$$

and define

$$\mathcal{H}_{\ell_1, \ell_2} = \{u \in \mathcal{A}_{\ell_1, \ell_2} \mid u_{x_1 y_1} + u_{x_2 y_2} + \cdots + u_{x_n y_n} = 0\}. \quad (1.10)$$

We proved in [32] that $\mathcal{H}_{\ell_1, \ell_2}$ forms an irreducible $sl(n, \mathbb{C})$ -submodule and

$$\mathcal{A}_{\ell_1, \ell_2} = \mathcal{H}_{\ell_1, \ell_2} \oplus (x_1 y_1 + x_2 y_2 + \cdots + x_n y_n) \mathcal{A}_{\ell_1-1, \ell_2-1}, \quad (1.11)$$

where $x_1 y_1 + x_2 y_2 + \cdots + x_n y_n$ is an $sl(n, \mathbb{C})$ -invariant. More importantly, an explicit basis for each $\mathcal{H}_{\ell_1, \ell_2}$ was constructed in [32].

Now we assume that $\mathcal{A}$ is the polynomial algebra in $x_1, \dots, x_7$. There exists an action of the simple Lie algebra $\mathcal{G}^{G_2}$ of type G_2 on $\mathcal{A}$, which keeps $x_1^2 + x_2 x_5 + x_3 x_6 + x_4 x_7$ invariant (e.g., cf. [32]). Again we denote by $\mathcal{A}_k$ the subspace of polynomials of degree k in $\mathcal{A}$ and define


$$\tilde{\mathcal{H}}_k = \{u \in \mathcal{A}_k \mid u_{x_1 x_1} + u_{x_2 x_5} + u_{x_3 x_6} + u_{x_4 x_7} = 0\}. \quad (1.12)$$

We showed in [32] that $\tilde{\mathcal{H}}_k$ forms an irreducible $\mathcal{G}^{G_2}$ -submodule and

$$\mathcal{A}_k = \tilde{\mathcal{H}}_k \oplus (x_1^2 + x_2 x_5 + x_3 x_6 + x_4 x_7) \mathcal{A}_{k-2}. \quad (1.13)$$

Moreover, an explicit basis for each $\tilde{\mathcal{H}}_k$ was constructed in [32]. Furthermore, Luo [24] generalized the result (1.5) to certain noncanonical polynomial representations of $o(n, \mathbb{C})$ and found a basis for the irreducible submodules by using our methods in [32]. The methods also enabled Cao [10] to obtain an explicit irreducible decomposition with respect to the real orthogonal Lie algebra for the space of polynomial solutions of the Navier equations in elasticity.

The Dynkin diagram of E_6 is as follows:



Denote by λ_i the i th fundamental weight of E_6 with respect to the above labeling. Let V be the 27-dimensional irreducible E_6 -module of highest weight λ_1 . Denote by $\mathcal{A}$ the polynomial algebra (equivalently, symmetric tensor) over V and by $\mathcal{A}_m$ the subspace of homogeneous polynomial with degree m . A *singular vector* in $\mathcal{A}$ is a weight vector annihilated by positive root vectors. The following is the main theorem of this paper:

Main Theorem: Any singular vector is a monomial in a linear singular vector x_1 of weight λ_1 , a quadratic singular vector ζ_1 of weight λ_6 and a cubic singular vector η of weight 0. In particular, η is the unique fundamental invariant and the following combinatorial identity holds:

$$(1-q)^{26} \sum_{m_1, m_2=0}^{\infty} (\dim V(m_1\lambda_1 + m_2\lambda_6)) q^{m_1+2m_2} = 1 + q + q^2. \quad (1.14)$$

Let $\mathcal{D}$ be the unique constant-coefficient fundamental invariant differential operator dual to η . Denote by $L(m_1, m_2)$ the E_6 -submodule generated by $x_1^{m_1} \zeta_1^{m_2}$. Then

$$\Phi_m = \{f \in \mathcal{A}_m \mid \mathcal{D}(f) = 0\} = \bigoplus_{i=0}^{\lfloor m/2 \rfloor} L(m-2i, i) \quad (1.15)$$

and

$$\mathcal{A}_m = \Phi_m \oplus \eta \mathcal{A}_{m-3}. \quad (1.16)$$

Note that the identities (1.1) and (1.14) are dimensional properties of irreducible submodules. Our identity (1.14) says that the dimensions of the irreducible E_6 -modules $V(m_1\lambda_1 + m_2\lambda_6)$ are correlated by the binomial coefficients of twenty-six. Moreover, η is the cubic polynomial invariant corresponding to the Dickson's invariant trilinear form (cf. [13]) and $\mathcal{D}$ is the invariant differential operator corresponding to the Dickson form. The equation (1.16) is exactly a cubic generalization of the quadratic ones in (1.5), (1.11) and (1.13). The fundamental difference is that our subspace Φ_m of homogeneous polynomial solutions is a sum of $\lfloor m/2 \rfloor + 1$ irreducible submodules.

In Section 2, we explicitly construct the 27-dimensional basic representation of E_6 in terms of differential operators via the root lattice construction of the E_7 simple Lie algebra. The proof of the main theorem is given in Section 3.


2 Basic Representation of E_6

In this section, we will explicitly construct the 27-dimensional basic irreducible representation of E_6 .

For convenience, we will use the notion

$$\overline{i, i+j} = \{i, i+1, i+2, \dots, i+j\} \quad (2.1)$$

for integer i and positive integer j throughout this paper. We start with the root lattice construction of the simple Lie algebra of type E_7 . As we all known, the Dynkin diagram of E_7 is as follows:



Let $\{\alpha_i \mid i \in \overline{1, 7}\}$ be the simple positive roots corresponding to the vertices in the diagram, and let Φ_{E_7} be the root system of E_7 . Set

$$Q_{E_7} = \sum_{i=1}^7 \mathbb{Z}\alpha_i, \quad (2.2)$$

the root lattice of type E_7 . Denote by $(\cdot, \cdot)$ the symmetric $\mathbb{Z}$ -bilinear form on Q_{E_7} such that

$$\Phi_{E_7} = \{\alpha \in Q_{E_7} \mid (\alpha, \alpha) = 2\}. \quad (2.3)$$

Define $F(\cdot, \cdot) : Q_{E_7} \times Q_{E_7} \rightarrow \{\pm 1\}$ by

$$F\left(\sum_{i=1}^7 k_i \alpha_i, \sum_{j=1}^7 l_j \alpha_j\right) = (-1)^{\sum_{i=1}^7 k_i l_i + \sum_{7 \geq i > j \geq 1} k_i l_j (\alpha_i, \alpha_j)}, \quad k_i, l_j \in \mathbb{Z}. \quad (2.4)$$

Then for $\alpha, \beta, \gamma \in Q_{E_7}$,

$$F(\alpha + \beta, \gamma) = F(\alpha, \gamma)F(\beta, \gamma), \quad F(\alpha, \beta + \gamma) = F(\alpha, \beta)F(\alpha, \gamma), \quad (2.5)$$

$$F(\alpha, \beta)F(\beta, \alpha)^{-1} = (-1)^{(\alpha, \beta)}, \quad F(\alpha, \alpha) = (-1)^{(\alpha, \alpha)/2}. \quad (2.6)$$

In particular,

$$F(\alpha, \beta) = -F(\beta, \alpha) \quad \text{if } \alpha, \beta, \alpha + \beta \in \Phi_{E_7}. \quad (2.7)$$

Denote

$$H_{E_7} = \sum_{i=1}^7 \mathbb{C}\alpha_i. \quad (2.8)$$

The simple Lie algebra of type E_7 is

$$\mathcal{G}^{E_7} = H_{E_7} \oplus \bigoplus_{\alpha \in \Phi_{E_7}} \mathbb{C}E_\alpha \quad (2.9)$$

with the Lie bracket $[\cdot, \cdot]$ determined by:

$$[H_{E_7}, H_{E_7}] = 0, \quad [h, E_\alpha] = (h, \alpha)E_\alpha, \quad [E_\alpha, E_{-\alpha}] = -\alpha, \quad (2.10)$$

$$[E_\alpha, E_\beta] = \begin{cases} 0 & \text{if } \alpha + \beta \notin \Phi_{E_7}, \\ F(\alpha, \beta)E_{\alpha+\beta} & \text{if } \alpha + \beta \in \Phi_{E_7}. \end{cases} \quad (2.11)$$

for $\alpha, \beta \in \Phi_{E_7}$ and $h \in H_{E_7}$.

Note that

$$Q_{E_6} = \sum_{i=1}^6 \mathbb{Z}\alpha_i \subset Q_{E_7} \quad (2.12)$$

is the root lattice of E_6 and

$$\Phi_{E_6} = Q_{E_6} \cap \Phi_{E_7} \quad (2.13)$$

is the root system of E_6 . Set

$$H_{E_6} = \sum_{i=1}^6 \mathbb{C}\alpha_i. \quad (2.14)$$

Then the subalgebra

$$\mathcal{G}^{E_6} = H_{E_6} \oplus \bigoplus_{\alpha \in \Phi_{E_6}} \mathbb{C}E_\alpha \quad (2.15)$$

of $\mathcal{G}^{E_7}$ is exactly the simple Lie algebra of $\mathcal{G}^{E_6}$. Denote by $\Phi_{E_6}^+$ the set of positive roots of E_6 and by $\Phi_{E_7}^+$ the set of positive roots of E_7 . The elements of $\Phi_{E_6}^+$ are:

$$\alpha_1 + 2\alpha_2 + 2\alpha_3 + 3\alpha_4 + 2\alpha_5 + \alpha_6, \quad (2.16)$$

$$\{\alpha_1 + \sum_{r=3}^j \alpha_r \mid j \in \overline{2, 6}\} \cup \{\sum_{r=i+1}^j \alpha_r \mid 2 \leq i < j \leq 6\}, \quad (2.17)$$

$$\{\sum_{s=2}^j \alpha_s + \sum_{t=4}^k \alpha_t \mid 2 \leq j < k \leq 6\} \quad (2.18)$$

and

$$\{\sum_{\iota=1}^i \alpha_\iota + \sum_{s=3}^j \alpha_s + \sum_{t=4}^k \alpha_t \mid 2 \leq i < j < k \leq 6\}. \quad (2.19)$$

Denote by $\bar{\Phi}_{E_7}^+$ the set of the following positive roots:

$$\alpha_1 + \sum_{r=3}^7 \alpha_r, \quad \alpha_3 + 2\alpha_4 + \alpha_5 + \sum_{i=1}^6 \alpha_i + \sum_{r=1}^7 \alpha_r, \quad (2.20)$$

$$\{2 \sum_{s=1}^6 \alpha_s - \alpha_1 + \alpha_4 - \alpha_6 + \sum_{r=i+1}^7 \alpha_r \mid i \in \overline{1, 6}\}, \quad \{\sum_{r=i+1}^7 \alpha_r \mid i \in \overline{2, 6}\}, \quad (2.21)$$

$$\{\sum_{s=2}^j \alpha_s + \sum_{t=4}^7 \alpha_t \mid j \in \overline{2, 6}\}, \quad \{\sum_{\iota=1}^i \alpha_\iota + \sum_{s=3}^j \alpha_s + \sum_{t=4}^7 \alpha_t \mid 2 \leq i < j \leq 6\}. \quad (2.22)$$

Then

$$\Phi_{E_7}^+ = \Phi_{E_6}^+ \cup \bar{\Phi}_{E_7}^+. \quad (2.23)$$

In particular,

$$V = \sum_{\beta \in \Phi_{E_7}^+} \mathbb{C}E_\beta \quad (2.24)$$

forms the 27-dimensional basic $\mathcal{G}^{E_6}$ -module of highest weight λ_1 with the representation $\text{ad}_{\mathcal{G}^{E_7}}$. Denote

$$x_1 = E_{\alpha_3+2\alpha_4+\alpha_5+\sum_{i=1}^6 \alpha_i+\sum_{r=1}^7 \alpha_r}, \quad x_2 = E_{2\sum_{s=1}^6 \alpha_s-\alpha_1+\alpha_4-\alpha_6+\sum_{r=3}^7 \alpha_r}, \quad (2.25)$$

$$x_3 = E_{2\sum_{s=1}^6 \alpha_s-\alpha_1+\alpha_4-\alpha_6+\sum_{r=4}^7 \alpha_r}, \quad x_4 = E_{2\sum_{s=1}^6 \alpha_s-\alpha_1-\alpha_6+\sum_{r=4}^7 \alpha_r} \quad (2.26)$$

$$x_5 = E_{2\sum_{s=1}^6 \alpha_s-\alpha_1+\alpha_4+\alpha_7}, \quad x_6 = E_{\sum_{t=1}^5 \alpha_t+\sum_{s=3}^6 \alpha_s+\sum_{t=4}^7 \alpha_t}, \quad (2.27)$$

$$x_7 = E_{\alpha_4+\sum_{s=3}^6 \alpha_s+\sum_{t=1}^7 \alpha_t}, \quad x_8 = E_{2\sum_{s=1}^6 \alpha_s-\alpha_1+\alpha_4-\alpha_6+\alpha_7}, \quad (2.28)$$

$$x_9 = E_{\sum_{s=3}^6 \alpha_s+\sum_{t=1}^7 \alpha_t}, \quad x_{10} = E_{\alpha_4+\sum_{s=3}^5 \alpha_s+\sum_{t=1}^7 \alpha_t}, \quad (2.29)$$

$$x_{11} = E_{\sum_{s=4}^6 \alpha_s+\sum_{t=1}^7 \alpha_t}, \quad x_{12} = E_{\sum_{s=3}^5 \alpha_s+\sum_{t=1}^7 \alpha_t}, \quad x_{13} = E_{\alpha_3+\alpha_4+\sum_{t=1}^7 \alpha_t}, \quad (2.30)$$

$$x_{14} = E_{\sum_{r=2}^6 \alpha_r+\sum_{s=4}^7 \alpha_s}, \quad x_{15} = E_{\alpha_4+\alpha_5+\sum_{t=1}^7 \alpha_t}, \quad x_{16} = E_{\alpha_4+\sum_{t=1}^7 \alpha_t}, \quad (2.31)$$

$$x_{17} = E_{\alpha_4+\alpha_5+\sum_{t=2}^7 \alpha_t}, \quad x_{18} = E_{\sum_{t=1}^7 \alpha_t}, \quad x_{19} = E_{\alpha_4+\sum_{t=2}^7 \alpha_t}, \quad (2.32)$$

$$x_{20} = E_{\alpha_1+\sum_{t=3}^7 \alpha_t}, \quad x_{21} = E_{\sum_{t=2}^7 \alpha_t}, \quad x_{22} = E_{\alpha_2+\sum_{t=4}^7 \alpha_t}, \quad x_{23} = E_{\sum_{t=3}^7 \alpha_t}, \quad (2.33)$$

$$x_{24} = E_{\sum_{t=4}^7 \alpha_t}, \quad x_{25} = E_{\sum_{t=5}^7 \alpha_t}, \quad x_{26} = E_{\alpha_6+\alpha_7}, \quad x_{27} = E_{\alpha_7}. \quad (2.34)$$

Then $\{x_i \mid i \in \overline{1, 27}\}$ forms a basis of V .

Under the above basis

$$E_{\alpha_1}|_V = -x_1\partial_{x_2} + x_{11}\partial_{x_{14}} + x_{15}\partial_{x_{17}} + x_{16}\partial_{x_{19}} + x_{18}\partial_{x_{21}} + x_{20}\partial_{x_{23}}, \quad (2.35)$$

$$E_{\alpha_2}|_V = -x_4\partial_{x_6} - x_5\partial_{x_7} - x_8\partial_{x_{10}} + x_{18}\partial_{x_{20}} + x_{21}\partial_{x_{23}} + x_{22}\partial_{x_{24}}, \quad (2.36)$$

$$E_{\alpha_3}|_V = -x_2\partial_{x_3} + x_9\partial_{x_{11}} + x_{12}\partial_{x_{15}} + x_{13}\partial_{x_{16}} + x_{21}\partial_{x_{22}} + x_{23}\partial_{x_{24}}, \quad (2.37)$$

$$E_{\alpha_4}|_V = -x_3\partial_{x_4} - x_7\partial_{x_9} - x_{10}\partial_{x_{12}} - x_{16}\partial_{x_{18}} - x_{19}\partial_{x_{21}} + x_{24}\partial_{x_{25}}, \quad (2.38)$$

$$E_{\alpha_5}|_V = -x_4\partial_{x_5} - x_6\partial_{x_7} - x_{12}\partial_{x_{13}} - x_{15}\partial_{x_{16}} - x_{17}\partial_{x_{19}} + x_{25}\partial_{x_{26}}, \quad (2.39)$$

$$E_{\alpha_6}|_V = -x_5\partial_{x_8} - x_7\partial_{x_{10}} - x_9\partial_{x_{12}} - x_{11}\partial_{x_{15}} - x_{14}\partial_{x_{17}} + x_{26}\partial_{x_{27}}, \quad (2.40)$$

$$E_{\alpha_1+\alpha_3}|_V = x_1\partial_{x_3} - x_9\partial_{x_{14}} - x_{12}\partial_{x_{17}} - x_{13}\partial_{x_{19}} + x_{18}\partial_{x_{22}} + x_{20}\partial_{x_{24}}, \quad (2.41)$$

$$E_{\alpha_2+\alpha_4}|_V = -x_3\partial_{x_6} + x_5\partial_{x_9} + x_8\partial_{x_{12}} + x_{16}\partial_{x_{20}} + x_{19}\partial_{x_{23}} + x_{22}\partial_{x_{25}}, \quad (2.42)$$

$$E_{\alpha_3+\alpha_4}|_V = x_2\partial_{x_4} + x_7\partial_{x_{11}} + x_{10}\partial_{x_{15}} - x_{13}\partial_{x_{18}} + x_{19}\partial_{x_{22}} + x_{23}\partial_{x_{25}}, \quad (2.43)$$

$$E_{\alpha_4+\alpha_5}|_V = x_3\partial_{x_5} - x_6\partial_{x_9} + x_{10}\partial_{x_{13}} - x_{15}\partial_{x_{18}} - x_{17}\partial_{x_{21}} + x_{24}\partial_{x_{26}}, \quad (2.44)$$

$$E_{\alpha_5+\alpha_6}|_V = x_4\partial_{x_8} + x_6\partial_{x_{10}} - x_9\partial_{x_{13}} - x_{11}\partial_{x_{16}} - x_{14}\partial_{x_{19}} + x_{25}\partial_{x_{27}}, \quad (2.45)$$

$$E_{\alpha_1+\alpha_3+\alpha_4}|_V = -x_1\partial_{x_4} - x_7\partial_{x_{14}} - x_{10}\partial_{x_{17}} + x_{13}\partial_{x_{21}} + x_{16}\partial_{x_{22}} + x_{20}\partial_{x_{25}}, \quad (2.46)$$

$$E_{\alpha_2+\alpha_3+\alpha_4}|_V = x_2\partial_{x_6} - x_5\partial_{x_{11}} - x_8\partial_{x_{15}} + x_{13}\partial_{x_{20}} - x_{19}\partial_{x_{24}} + x_{21}\partial_{x_{25}}, \quad (2.47)$$

$$E_{\alpha_2+\alpha_4+\alpha_5}|_V = x_3\partial_{x_7} + x_4\partial_{x_9} - x_8\partial_{x_{13}} + x_{15}\partial_{x_{20}} + x_{17}\partial_{x_{23}} + x_{22}\partial_{x_{26}}, \quad (2.48)$$

$$E_{\alpha_3+\alpha_4+\alpha_5}|_V = -x_2\partial_{x_5} + x_6\partial_{x_{11}} - x_{10}\partial_{x_{16}} - x_{12}\partial_{x_{18}} + x_{17}\partial_{x_{22}} + x_{23}\partial_{x_{26}}, \quad (2.49)$$

$$E_{\alpha_4+\alpha_5+\alpha_6}|_V = -x_3\partial_{x_8} + x_6\partial_{x_{12}} + x_7\partial_{x_{13}} - x_{11}\partial_{x_{18}} - x_{14}\partial_{x_{21}} + x_{24}\partial_{x_{27}}, \quad (2.50)$$

$$E_{\sum_{i=1}^4 \alpha_i}|_V = -x_1\partial_{x_6} + x_5\partial_{x_{14}} + x_8\partial_{x_{17}} - x_{13}\partial_{x_{23}} - x_{16}\partial_{x_{24}} + x_{18}\partial_{x_{25}}, \quad (2.51)$$

$$E_{\alpha_1+\sum_{i=3}^5 \alpha_i}|_V = x_1\partial_{x_5} - x_6\partial_{x_{14}} + x_{10}\partial_{x_{19}} + x_{12}\partial_{x_{21}} + x_{15}\partial_{x_{22}} + x_{20}\partial_{x_{26}}, \quad (2.52)$$

$$E_{\sum_{i=2}^5 \alpha_i}|_V = -x_2\partial_{x_7} - x_4\partial_{x_{11}} + x_8\partial_{x_{16}} + x_{12}\partial_{x_{20}} - x_{17}\partial_{x_{24}} + x_{21}\partial_{x_{26}}, \quad (2.53)$$

$$E_{\alpha_2+\sum_{i=4}^6 \alpha_i}|_V = -x_3\partial_{x_{10}} - x_4\partial_{x_{12}} - x_5\partial_{x_{13}} + x_{11}\partial_{x_{20}} + x_{14}\partial_{x_{23}} + x_{22}\partial_{x_{27}}, \quad (2.54)$$

$$E_{\sum_{i=3}^6 \alpha_i}|_V = x_2\partial_{x_8} - x_6\partial_{x_{15}} - x_7\partial_{x_{16}} - x_9\partial_{x_{18}} + x_{14}\partial_{x_{22}} + x_{23}\partial_{x_{27}}, \quad (2.55)$$

$$E_{\sum_{i=1}^5 \alpha_i}|_V = x_1\partial_{x_7} + x_4\partial_{x_{14}} - x_8\partial_{x_{19}} - x_{12}\partial_{x_{23}} - x_{15}\partial_{x_{24}} + x_{18}\partial_{x_{26}}, \quad (2.56)$$

$$E_{\alpha_1+\sum_{i=3}^6 \alpha_i}|_V = -x_1\partial_{x_8} + x_6\partial_{x_{17}} + x_7\partial_{x_{19}} + x_9\partial_{x_{21}} + x_{11}\partial_{x_{22}} + x_{20}\partial_{x_{27}}, \quad (2.57)$$

$$E_{\alpha_4+\sum_{i=2}^5 \alpha_i}|_V = x_2\partial_{x_9} - x_3\partial_{x_{11}} - x_8\partial_{x_{18}} + x_{10}\partial_{x_{20}} - x_{17}\partial_{x_{25}} + x_{19}\partial_{x_{26}}, \quad (2.58)$$

$$E_{\sum_{i=2}^6 \alpha_i}|_V = x_2\partial_{x_{10}} + x_4\partial_{x_{15}} + x_5\partial_{x_{16}} + x_9\partial_{x_{20}} - x_{14}\partial_{x_{24}} + x_{21}\partial_{x_{27}}, \quad (2.59)$$

$$E_{\alpha_4+\sum_{i=1}^5 \alpha_i}|_V = -x_1\partial_{x_9} + x_3\partial_{x_{14}} + x_8\partial_{x_{21}} - x_{10}\partial_{x_{23}} - x_{15}\partial_{x_{25}} + x_{16}\partial_{x_{26}}, \quad (2.60)$$

$$E_{\sum_{i=1}^6 \alpha_i}|_V = -x_1\partial_{x_{10}} - x_4\partial_{x_{17}} - x_5\partial_{x_{19}} - x_9\partial_{x_{23}} - x_{11}\partial_{x_{24}} + x_{18}\partial_{x_{27}}, \quad (2.61)$$

$$E_{\alpha_4+\sum_{i=2}^6 \alpha_i}|_V = -x_2\partial_{x_{12}} + x_3\partial_{x_{15}} - x_5\partial_{x_{18}} + x_7\partial_{x_{20}} - x_{14}\partial_{x_{25}} + x_{19}\partial_{x_{27}}, \quad (2.62)$$

$$E_{\alpha_3+\alpha_4+\sum_{i=1}^5 \alpha_i}|_V = x_1\partial_{x_{11}} - x_2\partial_{x_{14}} - x_8\partial_{x_{22}} + x_{10}\partial_{x_{24}} - x_{12}\partial_{x_{25}} + x_{13}\partial_{x_{26}}, \quad (2.63)$$

$$E_{\alpha_4+\sum_{i=1}^6 \alpha_i}|_V = x_1\partial_{x_{12}} - x_3\partial_{x_{17}} + x_5\partial_{x_{21}} - x_7\partial_{x_{23}} - x_{11}\partial_{x_{25}} + x_{16}\partial_{x_{27}}, \quad (2.64)$$

$$E_{\alpha_4+\alpha_5+\sum_{i=2}^6 \alpha_i}|_V = x_2\partial_{x_{13}} - x_3\partial_{x_{16}} - x_4\partial_{x_{18}} + x_6\partial_{x_{20}} - x_{14}\partial_{x_{26}} + x_{17}\partial_{x_{27}}, \quad (2.65)$$

$$E_{\alpha_3+\alpha_4+\sum_{i=1}^6 \alpha_i}|_V = -x_1\partial_{x_{15}} + x_2\partial_{x_{17}} - x_5\partial_{x_{22}} + x_7\partial_{x_{24}} - x_9\partial_{x_{25}} + x_{13}\partial_{x_{27}}, \quad (2.66)$$

$$E_{\alpha_4+\alpha_5+\sum_{i=1}^6 \alpha_i}|_V = -x_1\partial_{x_{13}} + x_3\partial_{x_{19}} + x_4\partial_{x_{21}} - x_6\partial_{x_{23}} - x_{11}\partial_{x_{26}} + x_{15}\partial_{x_{27}}, \quad (2.67)$$

$$E_{\sum_{r=3}^5 \alpha_r+\sum_{i=1}^6 \alpha_i}|_V = x_1\partial_{x_{16}} - x_2\partial_{x_{19}} - x_4\partial_{x_{22}} + x_6\partial_{x_{24}} - x_9\partial_{x_{26}} + x_{12}\partial_{x_{27}}, \quad (2.68)$$

$$E_{\alpha_4+\sum_{r=3}^5 \alpha_r+\sum_{i=1}^6 \alpha_i}|_V = x_1\partial_{x_{18}} - x_2\partial_{x_{21}} + x_3\partial_{x_{22}} - x_6\partial_{x_{25}} + x_7\partial_{x_{26}} - x_{10}\partial_{x_{27}}, \quad (2.69)$$

$$E_{\alpha_2+\alpha_4+\sum_{r=3}^5 \alpha_r+\sum_{i=1}^6 \alpha_i}|_V = x_1\partial_{x_{20}} - x_2\partial_{x_{23}} + x_3\partial_{x_{24}} - x_4\partial_{x_{25}} - x_5\partial_{x_{26}} - x_8\partial_{x_{27}}. \quad (2.70)$$

Recall that we also view α_i as the elements of $\mathcal{G}^{E_7}$ (cf. (2.8) and (2.9)). We write

$$[\alpha_j, x_i] = a_{i,j}x_i \quad \text{for } i \in \overline{1, 27}, j \in \overline{1, 6}. \quad (2.71)$$

Then the weight of x_i is $\sum_{j=1}^6 a_{i,j}\lambda_j$, where λ_j is the j th fundamental weight of $\mathcal{G}^{E_6}$. We calculate the following table:

Table 1

i	$a_{i,1}$	$a_{i,2}$	$a_{i,3}$	$a_{i,4}$	$a_{i,5}$	$a_{i,6}$	i	$a_{i,1}$	$a_{i,2}$	$a_{i,3}$	$a_{i,4}$	$a_{i,5}$	$a_{i,6}$
1	1	0	0	0	0	0	2	-1	0	1	0	0	0
3	0	0	-1	1	0	0	4	0	1	0	-1	1	0
5	0	1	0	0	-1	1	6	0	-1	0	0	1	0
7	0	-1	0	1	-1	1	8	0	1	0	0	0	-1
9	0	0	1	-1	0	1	10	0	-1	0	1	0	-1
11	1	0	-1	0	0	1	12	0	0	1	-1	1	-1
13	0	0	1	0	-1	0	14	-1	0	0	0	0	1
15	1	0	-1	0	1	-1	16	1	0	-1	1	-1	0
17	-1	0	0	0	1	-1	18	1	1	0	-1	0	0
19	-1	0	0	1	-1	0	20	1	-1	0	0	0	0
21	-1	1	1	-1	0	0	22	0	1	-1	0	0	0
23	-1	-1	1	0	0	0	24	0	-1	-1	1	0	0
25	0	0	0	-1	1	0	26	0	0	0	0	-1	1
27	0	0	0	0	0	-1							

In particular,

$$\alpha_j|_V = \sum_{i=1}^{27} a_{i,j} x_i \partial_{x_i} \quad \text{for } j \in \overline{1,6}. \quad (2.72)$$

Finally we have the representation formulas of the negative root vectors:

$$E_{-\alpha_1}|_V = x_2 \partial_{x_1} - x_{14} \partial_{x_{11}} - x_{17} \partial_{x_{15}} - x_{19} \partial_{x_{16}} - x_{21} \partial_{x_{18}} - x_{23} \partial_{x_{20}}, \quad (2.73)$$

$$E_{-\alpha_2}|_V = x_6 \partial_{x_4} + x_7 \partial_{x_5} + x_{10} \partial_{x_8} - x_{20} \partial_{x_{18}} - x_{23} \partial_{x_{21}} - x_{24} \partial_{x_{22}}, \quad (2.74)$$

$$E_{-\alpha_3}|_V = x_3 \partial_{x_2} - x_{11} \partial_{x_9} - x_{15} \partial_{x_{12}} - x_{16} \partial_{x_{13}} - x_{22} \partial_{x_{21}} - x_{24} \partial_{x_{23}}, \quad (2.75)$$

$$E_{-\alpha_4}|_V = x_4 \partial_{x_3} + x_9 \partial_{x_7} + x_{12} \partial_{x_{10}} + x_{18} \partial_{x_{16}} + x_{21} \partial_{x_{19}} - x_{25} \partial_{x_{24}}, \quad (2.76)$$

$$E_{-\alpha_5}|_V = x_5 \partial_{x_4} + x_7 \partial_{x_6} + x_{13} \partial_{x_{12}} + x_{16} \partial_{x_{15}} + x_{19} \partial_{x_{17}} - x_{26} \partial_{x_{25}}, \quad (2.77)$$

$$E_{-\alpha_6}|_V = x_8 \partial_{x_5} + x_{10} \partial_{x_7} + x_{12} \partial_{x_9} + x_{15} \partial_{x_{11}} + x_{17} \partial_{x_{14}} - x_{27} \partial_{x_{26}}, \quad (2.78)$$

$$E_{-\alpha_1-\alpha_3}|_V = -x_3 \partial_{x_1} + x_{14} \partial_{x_9} + x_{17} \partial_{x_{12}} + x_{19} \partial_{x_{13}} - x_{22} \partial_{x_{18}} - x_{24} \partial_{x_{20}}, \quad (2.79)$$

$$E_{-\alpha_2-\alpha_4}|_V = x_6 \partial_{x_3} - x_9 \partial_{x_5} - x_{12} \partial_{x_8} - x_{20} \partial_{x_{16}} - x_{23} \partial_{x_{19}} - x_{25} \partial_{x_{22}}, \quad (2.80)$$

$$E_{-\alpha_3-\alpha_4}|_V = -x_4 \partial_{x_2} - x_{11} \partial_{x_7} - x_{15} \partial_{x_{10}} + x_{18} \partial_{x_{13}} - x_{22} \partial_{x_{19}} - x_{25} \partial_{x_{23}}, \quad (2.81)$$

$$E_{-\alpha_4-\alpha_5}|_V = -x_5 \partial_{x_3} + x_9 \partial_{x_6} - x_{13} \partial_{x_{10}} + x_{18} \partial_{x_{15}} + x_{21} \partial_{x_{17}} - x_{26} \partial_{x_{24}}, \quad (2.82)$$

$$E_{-\alpha_5-\alpha_6}|_V = -x_8 \partial_{x_4} - x_{10} \partial_{x_6} + x_{13} \partial_{x_9} + x_{16} \partial_{x_{11}} + x_{19} \partial_{x_{14}} - x_{27} \partial_{x_{25}}, \quad (2.83)$$

$$E_{-\alpha_1-\alpha_3-\alpha_4}|_V = x_4 \partial_{x_1} + x_{14} \partial_{x_7} + x_{17} \partial_{x_{10}} - x_{21} \partial_{x_{13}} - x_{22} \partial_{x_{16}} - x_{25} \partial_{x_{20}}, \quad (2.84)$$

$$E_{-\alpha_2-\alpha_3-\alpha_4}|_V = -x_6 \partial_{x_2} + x_{11} \partial_{x_5} + x_{15} \partial_{x_8} - x_{20} \partial_{x_{13}} + x_{24} \partial_{x_{19}} - x_{25} \partial_{x_{21}}, \quad (2.85)$$

$$E_{-\alpha_2-\alpha_4-\alpha_5}|_V = -x_7\partial_{x_3} - x_9\partial_{x_4} + x_{13}\partial_{x_8} - x_{20}\partial_{x_{15}} - x_{23}\partial_{x_{17}} - x_{26}\partial_{x_{22}}, \quad (2.86)$$

$$E_{-\alpha_3-\alpha_4-\alpha_5}|_V = x_5\partial_{x_2} - x_{11}\partial_{x_6} + x_{16}\partial_{x_{10}} + x_{18}\partial_{x_{12}} - x_{22}\partial_{x_{17}} - x_{26}\partial_{x_{23}}, \quad (2.87)$$

$$E_{-\alpha_4-\alpha_5-\alpha_6}|_V = x_8\partial_{x_3} - x_{12}\partial_{x_6} - x_{13}\partial_{x_7} + x_{18}\partial_{x_{11}} + x_{21}\partial_{x_{14}} - x_{27}\partial_{x_{24}}, \quad (2.88)$$

$$E_{-\sum_{i=1}^4 \alpha_i}|_V = x_6\partial_{x_1} - x_{14}\partial_{x_5} - x_{17}\partial_{x_8} + x_{23}\partial_{x_{13}} + x_{24}\partial_{x_{16}} - x_{25}\partial_{x_{18}}, \quad (2.89)$$

$$E_{-\alpha_1-\sum_{i=3}^5 \alpha_i}|_V = -x_5\partial_{x_1} + x_{14}\partial_{x_6} - x_{19}\partial_{x_{10}} - x_{21}\partial_{x_{12}} - x_{22}\partial_{x_{15}} - x_{26}\partial_{x_{20}}, \quad (2.90)$$

$$E_{-\sum_{i=2}^5 \alpha_i}|_V = x_7\partial_{x_2} + x_{11}\partial_{x_4} - x_{16}\partial_{x_8} - x_{20}\partial_{x_{12}} + x_{24}\partial_{x_{17}} - x_{26}\partial_{x_{21}}, \quad (2.91)$$

$$E_{-\alpha_2-\sum_{i=4}^6 \alpha_i}|_V = x_{10}\partial_{x_3} + x_{12}\partial_{x_4} + x_{13}\partial_{x_5} - x_{20}\partial_{x_{11}} - x_{23}\partial_{x_{14}} - x_{27}\partial_{x_{22}}, \quad (2.92)$$

$$E_{-\sum_{i=3}^6 \alpha_i}|_V = -x_8\partial_{x_2} + x_{15}\partial_{x_6} + x_{16}\partial_{x_7} + x_{18}\partial_{x_9} - x_{22}\partial_{x_{14}} - x_{27}\partial_{x_{23}}, \quad (2.93)$$

$$E_{-\sum_{i=1}^5 \alpha_i}|_V = -x_7\partial_{x_1} - x_{14}\partial_{x_4} + x_{19}\partial_{x_8} + x_{23}\partial_{x_{12}} + x_{24}\partial_{x_{15}} - x_{26}\partial_{x_{18}}, \quad (2.94)$$

$$E_{-\alpha_1-\sum_{i=3}^6 \alpha_i}|_V = x_8\partial_{x_1} - x_{17}\partial_{x_6} - x_{19}\partial_{x_7} - x_{21}\partial_{x_9} - x_{22}\partial_{x_{11}} - x_{27}\partial_{x_{20}}, \quad (2.95)$$

$$E_{-\alpha_4-\sum_{i=2}^5 \alpha_i}|_V = -x_9\partial_{x_2} + x_{11}\partial_{x_3} + x_{18}\partial_{x_8} - x_{20}\partial_{x_{10}} + x_{25}\partial_{x_{17}} - x_{26}\partial_{x_{19}}, \quad (2.96)$$

$$E_{-\sum_{i=2}^6 \alpha_i}|_V = -x_{10}\partial_{x_2} - x_{15}\partial_{x_4} - x_{16}\partial_{x_5} - x_{20}\partial_{x_9} + x_{24}\partial_{x_{14}} - x_{27}\partial_{x_{21}}, \quad (2.97)$$

$$E_{-\alpha_4-\sum_{i=1}^5 \alpha_i}|_V = x_9\partial_{x_1} - x_{14}\partial_{x_3} - x_{21}\partial_{x_8} + x_{23}\partial_{x_{10}} + x_{25}\partial_{x_{15}} - x_{26}\partial_{x_{16}}, \quad (2.98)$$

$$E_{-\sum_{i=1}^6 \alpha_i}|_V = x_{10}\partial_{x_1} + x_{17}\partial_{x_4} + x_{19}\partial_{x_5} + x_{23}\partial_{x_9} + x_{24}\partial_{x_{11}} - x_{27}\partial_{x_{18}}, \quad (2.99)$$

$$E_{-\alpha_4-\sum_{i=2}^6 \alpha_i}|_V = x_{12}\partial_{x_2} - x_{15}\partial_{x_3} + x_{18}\partial_{x_5} - x_{20}\partial_{x_7} + x_{25}\partial_{x_{14}} - x_{27}\partial_{x_{19}}, \quad (2.100)$$

$$E_{-\alpha_3-\alpha_4-\sum_{i=1}^5 \alpha_i}|_V = -x_{11}\partial_{x_1} + x_{14}\partial_{x_2} + x_{22}\partial_{x_8} - x_{24}\partial_{x_{10}} + x_{25}\partial_{x_{12}} - x_{26}\partial_{x_{13}}, \quad (2.101)$$

$$E_{-\alpha_4-\sum_{i=1}^6 \alpha_i}|_V = -x_{12}\partial_{x_1} + x_{17}\partial_{x_3} - x_{21}\partial_{x_5} + x_{23}\partial_{x_7} + x_{25}\partial_{x_{11}} - x_{27}\partial_{x_{16}}, \quad (2.102)$$

$$E_{-\alpha_4-\alpha_5-\sum_{i=2}^6 \alpha_i}|_V = -x_{13}\partial_{x_2} + x_{16}\partial_{x_3} + x_{18}\partial_{x_4} - x_{20}\partial_{x_6} + x_{26}\partial_{x_{14}} - x_{27}\partial_{x_{17}}, \quad (2.103)$$

$$E_{-\alpha_3-\alpha_4-\sum_{i=1}^6 \alpha_i}|_V = x_{15}\partial_{x_1} - x_{17}\partial_{x_2} + x_{22}\partial_{x_5} - x_{24}\partial_{x_7} + x_{25}\partial_{x_9} - x_{27}\partial_{x_{13}}, \quad (2.104)$$

$$E_{-\alpha_4-\alpha_5-\sum_{i=1}^6 \alpha_i}|_V = x_{13}\partial_{x_1} - x_{19}\partial_{x_3} - x_{21}\partial_{x_4} + x_{23}\partial_{x_6} + x_{26}\partial_{x_{11}} - x_{27}\partial_{x_{15}}, \quad (2.105)$$

$$E_{-\sum_{r=3}^5 \alpha_r - \sum_{i=1}^6 \alpha_i}|_V = -x_{16}\partial_{x_1} + x_{19}\partial_{x_2} + x_{22}\partial_{x_4} - x_{24}\partial_{x_6} + x_{26}\partial_{x_9} - x_{27}\partial_{x_{12}}, \quad (2.106)$$

$$E_{-\alpha_4-\sum_{r=3}^5 \alpha_r - \sum_{i=1}^6 \alpha_i}|_V = x_{18}\partial_{x_1} - x_{21}\partial_{x_2} + x_{22}\partial_{x_3} - x_{25}\partial_{x_6} + x_{26}\partial_{x_7} - x_{27}\partial_{x_{10}}, \quad (2.107)$$

$$E_{-\alpha_2-\alpha_4-\sum_{r=3}^5 \alpha_r - \sum_{i=1}^6 \alpha_i}|_V = x_{20}\partial_{x_1} - x_{23}\partial_{x_2} + x_{24}\partial_{x_3} - x_{25}\partial_{x_4} + x_{26}\partial_{x_5} - x_{27}\partial_{x_8}. \quad (2.108)$$

3 Proof of the Main Theorem

Now $\mathcal{A} = \mathbb{C}[x_1, \dots, x_{27}]$ becomes a $\mathcal{G}^{E_6}$ -module via the differential operators in (2.35)-(2.108)

According to Table 1, we look for a singular vector of the form:

$$\zeta_1 = c_1 x_1 x_{14} + c_2 x_2 x_{11} + c_3 x_3 x_9 + c_4 x_4 x_7 + c_5 x_5 x_6. \quad (3.1)$$

By (2.35),

$$0 = E_{\alpha_1}(\zeta_1) = (c_1 - c_2)x_1 x_{11} \implies c_1 = c_2. \quad (3.2)$$

Moreover, (2.36) implies

$$0 = E_{\alpha_2(\zeta_1)} = -(c_4 + c_5)x_4 x_5 \implies c_5 = -c_4. \quad (3.3)$$

Furthermore, (2.37) gives

$$0 = E_{\alpha_3(\zeta_1)} = (c_2 - c_3)x_2 x_9 \implies c_2 = c_3. \quad (3.4)$$

In addition, (2.38) yields

$$0 = E_{\alpha_4(\zeta_1)} = -(c_3 + c_4)x_3 x_7 \implies c_4 = -c_3. \quad (3.5)$$

The last equation in (3.3) implies $E_{\alpha_5}(\zeta_1) = 0$ by (2.39). Besides, $E_{\alpha_6}(\zeta_1) = 0$ naturally holds by (3.1) and (2.40). Taking $c_1 = 1$, we have the singular vector

$$\zeta_1 = x_1 x_{14} + x_2 x_{11} + x_3 x_9 - x_4 x_7 + x_5 x_6 \quad (3.6)$$

of weight λ_6 .

According to (2.73)-(2.78), we set

$$\zeta_2 = E_{-\alpha_6}(\zeta_1) = x_1 x_{17} + x_2 x_{15} + x_3 x_{12} - x_4 x_{10} + x_6 x_8, \quad (3.7)$$

$$\zeta_3 = E_{-\alpha_5}(\zeta_2) = x_1 x_{19} + x_2 x_{16} + x_3 x_{13} - x_5 x_{10} + x_7 x_8, \quad (3.8)$$

$$\zeta_4 = E_{-\alpha_4}(\zeta_3) = x_1 x_{21} + x_2 x_{18} + x_4 x_{13} - x_5 x_{12} + x_8 x_9, \quad (3.9)$$

$$\zeta_5 = E_{-\alpha_3}(\zeta_4) = -x_1 x_{22} + x_3 x_{18} - x_4 x_{16} + x_5 x_{15} - x_8 x_{11}, \quad (3.10)$$

$$\zeta_6 = E_{-\alpha_2}(\zeta_4) = -x_1 x_{23} - x_2 x_{20} + x_6 x_{13} - x_7 x_{12} + x_9 x_{10}, \quad (3.11)$$

$$\zeta_7 = E_{-\alpha_3}(\zeta_6) = x_1 x_{24} - x_3 x_{20} - x_6 x_{16} + x_7 x_{15} - x_{10} x_{11}, \quad (3.12)$$

$$\zeta_8 = E_{-\alpha_1}(\zeta_5) = -x_2 x_{22} - x_3 x_{21} + x_4 x_{19} - x_5 x_{17} + x_8 x_{14}, \quad (3.13)$$

$$\zeta_9 = E_{-\alpha_4}(\zeta_7) = -x_1x_{25} - x_4x_{20} - x_6x_{18} + x_9x_{15} - x_{11}x_{12}, \quad (3.14)$$

$$\zeta_{10} = E_{-\alpha_1}(\zeta_7) = x_2x_{24} + x_3x_{23} + x_6x_{19} - x_7x_{17} + x_{10}x_{14}, \quad (3.15)$$

$$\zeta_{11} = -E_{-\alpha_5}(\zeta_9) = -x_1x_{26} + x_5x_{20} + x_7x_{18} - x_9x_{16} + x_{11}x_{13}, \quad (3.16)$$

$$\zeta_{12} = E_{-\alpha_4}(\zeta_{10}) = -x_2x_{25} + x_4x_{23} + x_6x_{21} - x_9x_{17} + x_{12}x_{14}, \quad (3.17)$$

$$\zeta_{13} = E_{-\alpha_3}(\zeta_{12}) = -x_3x_{25} - x_4x_{24} - x_6x_{22} + x_{11}x_{17} - x_{14}x_{15}, \quad (3.18)$$

$$\zeta_{14} = -E_{-\alpha_6}(\zeta_{11}) = -x_1x_{27} - x_8x_{20} - x_{10}x_{18} + x_{12}x_{16} - x_{13}x_{15}, \quad (3.19)$$

$$\zeta_{15} = -E_{-\alpha_5}(\zeta_{12}) = -x_2x_{26} - x_5x_{23} - x_7x_{21} + x_9x_{19} - x_{13}x_{14}. \quad (3.20)$$

Define a map $\iota : \overline{1, 27} \rightarrow \overline{1, 27}$ by

$$\iota(13) = 13, \quad \iota(14) = 14, \quad \iota(15) = 15, \quad (3.21)$$

$$\iota(i) = 28 - i \quad \text{for } i \in \overline{1, 27} \setminus \{13, 14, 15\}. \quad (3.22)$$

Let τ be an algebraic automorphism of $\mathcal{A}$ determined by

$$\tau(x_i) = x_{\iota(i)} \quad \text{for } i \in \overline{1, 27}. \quad (3.23)$$

Now we set

$$\zeta_i = \tau(\zeta_{28-i}) \quad \text{for } i \in \overline{16, 27}. \quad (3.24)$$

It can be verified that

$$\bar{V} = \sum_{r=1}^{27} \mathbb{C}\zeta_r \quad (3.25)$$

an irreducible $\mathcal{G}^{E_6}$ -submodule and $\{\zeta_r \mid r \in \overline{1, 27}\}$ forms a basis of $\bar{V}$.

From the Dynkin diagram of E_6 , we have the following automorphism of Q_{E_6} :

$$\sigma\left(\sum_{i=1}^6 k_i \alpha_i\right) = k_6 \alpha_1 + k_2 \alpha_2 + k_5 \alpha_3 + k_4 \alpha_4 + k_3 \alpha_5 + k_1 \alpha_6 \quad (3.26)$$

for $\sum_{i=1}^6 k_i \alpha_i \in Q_{E_6}$. Let ν be an associative algebra homomorphism of the associative algebra

$$\mathbb{A} = \sum_{i_1, \dots, i_{27}=0}^{\infty} \mathcal{A} \partial_{x_1}^{i_1} \cdots \partial_{x_{27}}^{i_{27}} \quad (3.27)$$

of differential operators to itself determined by

$$\nu(x_i) = \zeta_i, \quad \nu(\partial_{x_i}) = \partial_{\zeta_i} \quad \text{for } i \in \overline{1, 27}. \quad (3.28)$$

It can be proved that

$$E_\alpha|_{\bar{V}} = \nu(E_{\sigma(\alpha)}|_V) \quad \text{for } \alpha \in \Phi_{E_6}^+. \quad (3.29)$$

Moreover,

$$\alpha_j|_{\bar{V}} = \sum_{i=1}^{27} b_{i,j} \zeta_i \partial_{\zeta_i}, \quad (3.30)$$

where

$$b_{i,1} = a_{i,6}, \quad b_{i,3} = a_{i,5}, \quad b_{i,2} = a_{i,2}, \quad b_{i,4} = a_{i,4}. \quad (3.31)$$

Thus we have the following table:

Table 2

i	$b_{i,1}$	$b_{i,2}$	$b_{i,3}$	$b_{i,4}$	$b_{i,5}$	$b_{i,6}$	i	$b_{i,1}$	$b_{i,2}$	$b_{i,3}$	$b_{i,4}$	$b_{i,5}$	$b_{i,6}$
1	0	0	0	0	0	1	2	0	0	0	0	1	-1
3	0	0	0	1	-1	0	4	0	1	1	-1	0	0
5	1	1	-1	0	0	0	6	0	-1	1	0	0	0
7	1	-1	-1	1	0	0	8	-1	1	0	0	0	0
9	1	0	0	-1	1	0	10	-1	-1	0	1	0	0
11	1	0	0	0	-1	1	12	-1	0	1	-1	1	0
13	0	0	-1	0	1	0	14	1	0	0	0	0	-1
15	-1	0	1	0	-1	1	16	0	0	-1	1	-1	1
17	-1	0	1	0	0	-1	18	0	1	0	-1	0	1
19	0	0	-1	1	0	-1	20	0	-1	0	0	0	1
21	0	1	0	-1	1	-1	22	0	1	0	0	-1	0
23	0	-1	0	0	1	-1	24	0	-1	0	1	-1	0
25	0	0	1	-1	0	0	26	1	0	-1	0	0	0
27	-1	0	0	0	0	0							

According to Table 1 and Table 2, we look for an invariant of the form

$$\eta = \sum_{i=1}^{12} (d_i x_i \zeta_{28-i} + d_{28-i} x_{28-i} \zeta_i) + d_{13} x_{13} \zeta_{13} + d_{14} x_{14} \zeta_{14} + d_{15} x_{15} \zeta_{15}, \quad (3.32)$$

where $d_i \in \mathbb{C}$. By (2.35), (2.40) and (3.29), we have

$$\begin{aligned} 0 &= E_{\alpha_1}(\eta) \\ &= -d_2 x_1 \zeta_{26} + d_{14} x_{11} \zeta_{14} + d_{17} x_{15} \zeta_{11} + d_{19} x_{16} \zeta_9 + d_{21} x_{18} \zeta_7 + d_{23} x_{20} \zeta_5 \\ &\quad - d_{20} x_{20} \zeta_5 - d_{18} x_{18} \zeta_7 - d_{16} x_{16} \zeta_9 - d_{15} x_{15} \zeta_{11} - d_{11} x_{11} \zeta_{14} + d_1 x_1 \zeta_{26}, \end{aligned} \quad (3.33)$$

$$\begin{aligned} 0 &= E_{\alpha_6}(\eta) \\ &= -d_8 x_5 \zeta_{20} - d_{10} x_7 \zeta_{18} - d_{12} x_9 \zeta_{16} - d_{15} x_{11} \zeta_{15} - d_{17} x_{14} \zeta_{11} + d_{27} x_{26} \zeta_1 \\ &\quad - d_{26} x_{26} \zeta_1 + d_{14} x_{14} \zeta_{11} + d_{11} x_{11} \zeta_{15} + d_9 x_9 \zeta_{16} + d_7 x_7 \zeta_{18} + d_5 x_5 \zeta_{20}. \end{aligned} \quad (3.34)$$

So we take

$$d_2 = d_1, \quad d_{14} = d_{11}, \quad d_{17} = d_{15}, \quad d_{19} = d_{16}, \quad d_{21} = d_{18}, \quad d_{23} = d_{20}, \quad (3.35)$$

$$d_8 = d_7, \quad d_{10} = d_7, \quad d_{12} = d_9, \quad d_{15} = d_{11}, \quad d_{17} = d_{14}, \quad d_{27} = d_{26}. \quad (3.36)$$

Moreover, (2.37), (2.39) and (3.29) imply

$$\begin{aligned} 0 &= E_{\alpha_3}(\eta) \\ &= -d_3x_2\zeta_{25} + d_{11}x_9\zeta_{17} + d_{15}x_{12}\zeta_{15} + d_{16}x_{13}\zeta_{12} + d_{22}x_{21}\zeta_6 + d_{24}x_{23}\zeta_4 \\ &\quad -d_{23}x_{23}\zeta_4 - d_{21}x_{21}\zeta_6 - d_{13}x_{13}\zeta_{12} - d_{12}x_{12}\zeta_{15} - d_9x_9\zeta_{17} + d_2x_2\zeta_{25}, \end{aligned} \quad (3.37)$$

$$\begin{aligned} 0 &= E_{\alpha_5}(\eta) \\ &= -d_5x_4\zeta_{23} - d_7x_6\zeta_{21} - d_{13}x_{12}\zeta_{13} - d_{16}x_{15}\zeta_{12} - d_{19}x_{17}\zeta_9 + d_{26}x_{25}\zeta_2 \\ &\quad -d_{25}x_{25}\zeta_2 + d_{17}x_{17}\zeta_9 + d_{15}x_{15}\zeta_{12} + d_{12}x_{12}\zeta_{13} + d_6x_6\zeta_{21} + d_4x_4\zeta_{23}. \end{aligned} \quad (3.38)$$

Hence we get

$$d_3 = d_2, \quad d_{11} = d_9, \quad d_{15} = d_{12}, \quad d_{16} = d_{13}, \quad d_{22} = d_{21}, \quad d_{24} = d_{23}, \quad (3.39)$$

$$d_5 = d_4, \quad d_7 = d_6, \quad d_{13} = d_{12}, \quad d_{16} = d_{15}, \quad d_{19} = d_{17}, \quad d_{26} = d_{25}. \quad (3.40)$$

Furthermore, (2.36), (2.38) and (3.29) yield

$$\begin{aligned} 0 &= E_{\alpha_2}(\eta) \\ &= -d_6x_4\zeta_{22} - d_{22}x_{22}\zeta_4 - d_7x_5\zeta_{21} - d_{21}x_{21}\zeta_5 - d_{10}x_8\zeta_{18} - d_{18}x_{18}\zeta_8 \\ &\quad +d_{20}x_{18}\zeta_8 + d_8x_8\zeta_{18} + d_{23}x_{21}\zeta_5 + d_5x_5\zeta_{21} + d_{24}x_{22}\zeta_4 + d_4x_4\zeta_{22}, \end{aligned} \quad (3.41)$$

$$\begin{aligned} 0 &= E_{\alpha_4}(\eta) \\ &= -d_4x_3\zeta_{24} - d_{24}x_{24}\zeta_3 - d_9x_7\zeta_{19} - d_{19}x_{19}\zeta_7 - d_{12}x_{10}\zeta_{16} - d_{16}x_{16}\zeta_{10} \\ &\quad -d_{18}x_{16}\zeta_{10} - d_{10}x_{10}\zeta_{16} - d_{21}x_{19}\zeta_7 - d_7x_7\zeta_{19} + d_{25}x_{24}\zeta_3 + d_3x_3\zeta_{24}. \end{aligned} \quad (3.42)$$

Thus we obtain

$$d_6 = d_4, \quad d_{24} = d_{22}, \quad d_7 = d_5, \quad d_{23} = d_{21}, \quad d_{10} = d_8, \quad d_{20} = d_{18}, \quad (3.43)$$

$$d_4 = d_3, \quad d_{25} = d_{24}, \quad d_9 = -d_7, \quad d_{21} = -d_{19}, \quad d_{12} = -d_{10}, \quad d_{18} = -d_{16}. \quad (3.44)$$

By (3.35), (3.36), (3.39), (3.40), (3.42) and (3.43), we have

$$\begin{aligned} d_1 &= d_2 = d_3 = d_4 = d_5 = d_6 = d_7 = d_8 = -d_9 = d_{10} = -d_{11} \\ &= -d_{12} = -d_{13} = -d_{14} = -d_{15} = -d_{16} = -d_{17} = d_{18} \\ &= -d_{19} = d_{20} = d_{21} = d_{22} = d_{23} = d_{24} = d_{25} = d_{26} = d_{27}. \end{aligned} \quad (3.45)$$

Therefore, we have the following invariant

$$\eta = \sum_{i=1}^8 (x_i \zeta_{28-i} + x_{28-i} \zeta_i) + x_{10} \zeta_{18} - \sum_{r=9,11,12} (x_r \zeta_{28-r} + x_{28-r} \zeta_r) - \sum_{s=13}^{15} x_s \zeta_s. \quad (3.46)$$

According to (3.6)-(3.24),

$$\begin{aligned} \eta = & 3[(x_1 x_{14} + x_2 x_{11} + x_3 x_9) x_{27} + (x_1 x_{17} + x_2 x_{15} + x_3 x_{12}) x_{26} + (x_1 x_{19} + x_2 x_{16} \\ & + x_3 x_{13}) x_{25} + (x_4 x_{13} - x_5 x_{12} + x_8 x_9) x_{24} - (x_4 x_{16} - x_5 x_{15} + x_8 x_{11}) x_{23} + (x_6 x_{13} \\ & - x_7 x_{12} + x_9 x_{10}) x_{22} + (x_7 x_{15} + x_6 x_{16} - x_{10} x_{11}) x_{21} + (x_4 x_{19} - x_5 x_{17} + x_8 x_{14}) x_{20} \\ & + (x_6 x_{18} - x_9 x_{15} + x_{11} x_{12}) x_{19} + (x_{10} x_{14} - x_7 x_{17}) x_{18} + (x_9 x_{16} - x_{11} x_{13}) x_{17} \\ & - x_{12} x_{14} x_{16} + x_{14} x_{15} x_{13}] + (x_4 x_7 - x_5 x_6) x_{27} + (x_4 x_{10} - x_6 x_8) x_{26} + (x_5 x_{10} \\ & - x_7 x_8) x_{25} + (x_1 x_{21} + x_2 x_{18}) x_{24} + (x_3 x_{18} - x_1 x_{22}) x_{23} - (x_2 x_{22} + x_3 x_{21}) x_{20}. \end{aligned} \quad (3.47)$$

Lemma 3.1. *Any homogeneous singular vector in $\mathcal{A}$ is a monomial in x_1 , ζ_1 and η .*

Proof. Note that

$$x_1 x_{14} = \zeta_1 - x_2 x_{11} - x_3 x_9 + x_4 x_7 - x_5 x_6 \quad (3.48)$$

$$x_1 x_{17} = \zeta_2 - x_2 x_{15} - x_3 x_{12} + x_4 x_{10} - x_6 x_8, \quad (3.49)$$

$$x_1 x_{19} = \zeta_3 - x_2 x_{16} - x_3 x_{13} + x_5 x_{10} - x_7 x_8, \quad (3.50)$$

$$x_1 x_{21} = \zeta_4 - x_2 x_{18} - x_4 x_{13} + x_5 x_{12} - x_8 x_9, \quad (3.51)$$

$$x_1 x_{22} = -\zeta_5 - x_3 x_{18} - x_4 x_{16} + x_5 x_{15} - x_8 x_{11}, \quad (3.52)$$

$$x_1 x_{23} = -\zeta_6 - x_2 x_{20} + x_6 x_{13} - x_7 x_{12} + x_9 x_{10}, \quad (3.53)$$

$$x_1 x_{24} = \zeta_7 + x_3 x_{20} + x_6 x_{16} - x_7 x_{15} + x_{10} x_{11}, \quad (3.54)$$

$$x_1 x_{25} = -\zeta_9 - x_4 x_{20} - x_6 x_{18} + x_9 x_{15} - x_{11} x_{12}, \quad (3.55)$$

$$x_1 x_{26} = -\zeta_{11} + x_5 x_{20} + x_7 x_{18} - x_9 x_{16} + x_{11} x_{13} \quad (3.56)$$

by (3.6)-(3.12), (3.14) and (3.16). Moreover, (3.47) can be written as

$$\begin{aligned} & (3x_1 x_{14} + 3x_2 x_{11} + 3x_3 x_9 + x_4 x_7 - x_5 x_6) x_{27} \\ = & \eta - 3[(x_1 x_{17} + x_2 x_{15} + x_3 x_{12}) x_{26} + (x_1 x_{19} + x_2 x_{16} + x_3 x_{13}) x_{25} + (x_4 x_{13} \\ & - x_5 x_{12} + x_8 x_9) x_{24} - (x_4 x_{16} - x_5 x_{15} + x_8 x_{11}) x_{23} + (x_6 x_{13} - x_7 x_{12} + x_9 x_{10}) x_{22} \\ & + (x_7 x_{15} + x_6 x_{16} - x_{10} x_{11}) x_{21} + (x_4 x_{19} - x_5 x_{17} + x_8 x_{14}) x_{20} + (x_6 x_{18} \\ & - x_9 x_{15} + x_{11} x_{12}) x_{19} + (x_{10} x_{14} - x_7 x_{17}) x_{18} + (x_9 x_{16} - x_{11} x_{13}) x_{17} \\ & - x_{12} x_{14} x_{16} + x_{14} x_{15} x_{13}] - (x_4 x_{10} - x_6 x_8) x_{26} - (x_5 x_{10} - x_7 x_8) x_{25} \\ & - (x_1 x_{21} + x_2 x_{18}) x_{24} - (x_3 x_{18} - x_1 x_{22}) x_{23} + (x_2 x_{22} + x_3 x_{21}) x_{20}. \end{aligned} \quad (3.57)$$

Let f be any homogenous singular vector in $\mathcal{A}$. According to the above equations, f can be written as a rational function f_1 in

$$\{x_i, \zeta_r, \eta \mid i \in \{\overline{1, 13}, 15, 16, 18, 20\}; r \in \{\overline{1, 7}, 9, 11\}\}. \quad (3.58)$$

By (2.63)-(2.70), (3.28) and (3.29),

$$0 = E_{\alpha_3 + \alpha_4 + \sum_{i=1}^5 \alpha_i}(f_1) = x_1 \partial_{x_{11}}(f_1), \quad (3.59)$$

$$0 = E_{\alpha_4 + \sum_{i=1}^6 \alpha_i}(f_1) = x_1 \partial_{x_{12}}(f_1), \quad (3.60)$$

$$0 = E_{\alpha_4 + \alpha_5 + \sum_{i=2}^6 \alpha_i}(f_1) = x_2 \partial_{x_{13}}(f_1) + \zeta_1 \partial_{\zeta_{11}}(f_1), \quad (3.61)$$

$$0 = E_{\alpha_3 + \alpha_4 + \sum_{i=1}^6 \alpha_i}(f_1) = -x_1 \partial_{x_{15}}(f_1), \quad (3.62)$$

$$0 = E_{\alpha_4 + \alpha_5 + \sum_{i=1}^6 \alpha_i}|_V = -x_1 \partial_{x_{13}}(f_1), \quad (3.63)$$

$$0 = E_{\sum_{r=3}^5 \alpha_r + \sum_{i=1}^6 \alpha_i}|_V = x_1 \partial_{x_{16}}(f_1), \quad (3.64)$$

$$0 = E_{\alpha_4 + \sum_{r=3}^5 \alpha_r + \sum_{i=1}^6 \alpha_i}|_V = x_1 \partial_{x_{18}}(f_1), \quad (3.65)$$

$$0 = E_{\alpha_2 + \alpha_4 + \sum_{r=3}^5 \alpha_r + \sum_{i=1}^6 \alpha_i}(f_1) = x_1 \partial_{x_{20}}(f_1). \quad (3.66)$$

So f_1 is independent of $x_{11}, x_{12}, x_{13}, x_{15}, x_{16}, x_{18}, x_{20}$ and ζ_{11} , that is, f_1 is a rational function in

$$\{x_i, \zeta_r, \eta \mid i \in \overline{1, 10}; r \in \{\overline{1, 7}, 9\}\}. \quad (3.67)$$

Next (2.56)-(2.62), (3.28) and (3.29) imply that

$$0 = E_{\sum_{i=1}^5 \alpha_i}(f_1) = x_1 \partial_{x_7}(f_1), \quad (3.68)$$

$$0 = E_{\alpha_1 + \sum_{i=3}^6 \alpha_i}(f_1) = -x_1 \partial_{x_8}(f_1), \quad (3.69)$$

$$0 = E_{\alpha_4 + \sum_{i=2}^5 \alpha_i}(f_1) = x_2 \partial_{x_9}(f_1) + \zeta_2 \partial_{\zeta_9}(f_1), \quad (3.70)$$

$$0 = E_{\sum_{i=2}^6 \alpha_i}(f_1) = x_2 \partial_{x_{10}}(f_1) + \zeta_1 \partial_{\zeta_7}(f_1), \quad (3.71)$$

$$0 = E_{\alpha_4 + \sum_{i=1}^5 \alpha_i}(f_1) = -x_1 \partial_{x_9}(f_1), \quad (3.72)$$

$$0 = E_{\sum_{i=1}^6 \alpha_i}(f_1) = -x_1 \partial_{x_{10}}(f_1). \quad (3.73)$$

Hence f_1 is independent of $x_7, x_8, x_9, x_{10}, \zeta_7$ and ζ_9 , that is, f_1 is a rational function in

$$\{x_i, \zeta_r, \eta \mid i, r \in \overline{1, 6}\}. \quad (3.74)$$

Now (2.41), (2.45)-(2.50), (2.52), (2.55), (3.28) and (3.29) give that

$$0 = E_{\alpha_1 + \alpha_3}(f_1) = x_1 \partial_{x_3}(f_1), \quad (3.75)$$

$$0 = E_{\alpha_5 + \alpha_6}(f_1) = \zeta_1 \partial_{\zeta_3}(f_1), \quad (3.76)$$

$$0 = E_{\alpha_1 + \alpha_3 + \alpha_4}(f_1) = -x_1 \partial_{x_4}(f_1), \quad (3.77)$$

$$0 = E_{\alpha_2 + \alpha_3 + \alpha_4}(f_1) = x_2 \partial_{x_6}(f_1), \quad (3.78)$$

$$0 = E_{\alpha_2 + \alpha_4 + \alpha_5}|_V = \zeta_2 \partial_{\zeta_6}(f_1), \quad (3.79)$$

$$0 = E_{\alpha_3 + \alpha_4 + \alpha_5}(f_1) = -x_2 \partial_{x_5}(f_1) - \zeta_2 \partial_{\zeta_5}(f_1), \quad (3.80)$$

$$0 = E_{\alpha_4 + \alpha_5 + \alpha_6}(f_1) = -\zeta_1 \partial_{\zeta_4}(f_1), \quad (3.81)$$

$$0 = E_{\alpha_1 + \sum_{i=3}^5 \alpha_i}(f_1) = x_1 \partial_{x_5}(f_1). \quad (3.82)$$

Thus f_1 is independent of $\{x_i, \zeta_i \mid i \in \overline{3, 6}\}$, that is, f_1 is a rational function in $\{x_1, x_2, \zeta_1, \zeta_2, \eta\}$. Finally, (2.35), (2.40), (3.28) and (3.29) yield

$$0 = E_{\alpha_1}(f_1) = -x_1 \partial_{x_2}(f_1), \quad 0 = E_{\alpha_6}(f_1) = -\zeta_1 \partial_{\zeta_2}(f_1). \quad (3.83)$$

Therefore, f_1 is independent of x_2 and ζ_2 , that is, $f = f_1$ is a rational function in x_1 , ζ_1 and η . By (3.48) and (3.57), it must be a polynomial in x_1 , ζ_1 and η . Recall that the weights of x_1 , ζ_1 and η are λ_1 , λ_6 and 0, respectively. The homogeneity of f implies that it must be a monomial in x_1 , ζ_1 and η . $\square$

Let $L(m_1, m_2, m_3)$ be the $\mathcal{G}^{E_6}$ -submodule generated by $x_1^{m_1} \zeta_1^{m_2} \eta^{m_3}$. Then $L(m_1, m_2, m_3)$ is a finite-dimensional irreducible submodule of highest weight $m_1 \lambda_1 + m_2 \lambda_6$. By the Weyl's theorem of completely reducibility and the above lemma, we have

$$\mathcal{A} = \sum_{m_1, m_2, m_3=0}^{\infty} L(m_1, m_2, m_3). \quad (3.84)$$

Recall we denote by $V(\lambda)$ the finite-dimensional irreducible module of highest weight λ . The above equation implies

$$\frac{1}{(1-q)^{27}} = \frac{1}{1-q^3} \sum_{m_1, m_2=0}^{\infty} (\dim V(m_1 \lambda_1 + m_2 \lambda_6)) q^{m_1 + 2m_2}. \quad (3.85)$$

Equivalently, we have:

Lemma 3.2. *The following dimensional property of irreducible $\mathcal{G}^{E_6}$ -modules holds:*

$$(1-q)^{26} \sum_{m_1, m_2=0}^{\infty} (\dim V(m_1 \lambda_1 + m_2 \lambda_6)) q^{m_1 + 2m_2} = 1 + q + q^2. \quad (3.86)$$

Set

$$W = \sum_{i=1}^{27} \mathbb{C} \partial_{x_i}. \quad (3.87)$$

Then W is isomorphic to the module of linear functions on V via $\partial_{x_i}(x_j) = \delta_{i,j}$. Indeed, the linear map determined by $\partial_{x_i} \mapsto \zeta_{\iota(i)}$ (cf. (3.21), (3.22)) is a $\mathcal{G}^{E_6}$ -module isomorphism. We define a linear map $\mathfrak{S} : \mathcal{A} \rightarrow \mathbb{C}[\partial_{x_1}, \dots, \partial_{x_{27}}]$ by

$$\mathfrak{S}(x_1^{\alpha_1} x_2^{\alpha_2} \dots x_{27}^{\alpha_{27}}) = \partial_{x_1}^{\alpha_1} \partial_{x_2}^{\alpha_2} \dots \partial_{x_{27}}^{\alpha_{27}}. \quad (3.89)$$

Set

$$\mathcal{D} = \mathfrak{S}(\eta), \quad \mathcal{D}_1 = \sum_{i=1}^{27} x_i \partial_{x_i}, \quad \mathcal{D}_2 = \sum_{i=1}^{27} \zeta_i \mathfrak{S}(\zeta_i). \quad (3.89)$$

Then $\mathcal{D}$, $\mathcal{D}_1$ and $\mathcal{D}_2$ are invariant differential operators, that is,

$$(\mathcal{D}\xi)|_{\mathcal{A}} = (\xi\mathcal{D})|_{\mathcal{A}}, \quad (\mathcal{D}_r\xi)|_{\mathcal{A}} = (\xi\mathcal{D}_r)|_{\mathcal{A}} \quad \text{for } \xi \in \mathcal{G}^{E_6}. \quad (3.90)$$

Note that Lemma 3.1 implies

$$V^2 = L(2, 0, 0) + L(0, 1, 0). \quad (3.91)$$

Symmetrically,

$$W^2 = L'(0, 2, 0) + L'(1, 0, 0), \quad (3.92)$$

where $L'(0, 2, 0)$ is a module generated by the highest weight vector ∂_{27}^2 with weight $2\lambda_6$ and $L'(1, 0, 0)$ is a module generated by the highest weight vector $\mathfrak{S}(\zeta_{27})$ with weight λ_1 . Thus the subspace of invariants (the trivial submodule) in V^2W^2 is two-dimensional. The trivial submodule of $L(0, 1, 0)L'(1, 0, 0)$ is $\mathbb{C}\mathcal{D}_2$. In $L(2, 0, 0)L'(0, 2, 0)$, there exists an invariant $\mathcal{D}_3$ with a term $x_1^2 \partial_{x_1}^2$. So any invariant in V^2W^2 must be in $\mathbb{C}\mathcal{D}_2 + \mathbb{C}\mathcal{D}_3$. In particular, the invariant differential operator

$$[\mathcal{D}, \eta] = \mathcal{D}\eta - \eta\mathcal{D} = b_0 + b_1\mathcal{D}_1 + b_2\mathcal{D}_2 + b_3\mathcal{D}_3 \quad (3.93)$$

for some $b_s \in \mathbb{C}$. According to (3.47), η does not contain x_1^2 . So $b_3 = 0$. Moreover, (3.47) also implies $b_0 = 11$.

According to (3.57), the coefficient of $x_{27}\partial_{x_{27}}$ in $[\mathcal{D}_0, \eta]$ must be 11, which implies $b_1 = 11$. Observe that there exists a unique monomial in η containing x_1x_{14} , which is $3x_1x_{14}x_{27}$. Thus the coefficient of $x_1x_{14}\partial_{x_1}\partial_{x_{14}}$ in $[\mathcal{D}, \eta]$ must be 9, that is, $b_2 = 9$. So we have:

Lemma 3.3. *As operators on $\mathcal{A}$,*

$$[\mathcal{D}, \eta] = 111 + 11\mathcal{D}_1 + 9\mathcal{D}_2. \quad (3.94)$$

Let m_1 and m_2 be nonnegative integers. If $\mathcal{D}(x_1^{m_1}\zeta_1^{m_2}) \neq 0$, then it is also a singular of degree $m_1 + 2m_2 - 3$ with the same weight $m_1\lambda_1 + m_2\lambda_6$. But Lemma 3.1 implies that any singular vector with weight $m_1\lambda_1 + m_2\lambda_6$ must has degree $\geq m_1 + 2m_2$. This leads a contradiction. Thus

$$\mathcal{D}(x_1^{m_1}\zeta_1^{m_2}) = 0 \quad \text{for } m_1, m_2 \in \mathbb{N}. \quad (3.95)$$

Moreover, (3.90) implies

$$\mathcal{D}(L(m_1, m_2, 0)) = \{0\} \quad \text{for } m_1, m_2 \in \mathbb{N}. \quad (3.96)$$

Since $\mathcal{D}_2(x_1^{m_1}\zeta_1^{m_2})$ is also a singular vector of degree $m_1 + 2m_2$ with the same weight $m_1\lambda_1 + m_2\lambda_6$, we have

$$\mathcal{D}_2(x_1^{m_1}\zeta_1^{m_2}) = cx_1^{m_1}\zeta_1^{m_2} \quad (3.97)$$

for some $c \in \mathbb{C}$. Let

$$x_i = 0 \quad \text{for } 1, 14 \neq i \in \overline{1, 27} \quad (3.98)$$

in (3.97) and we get

$$\begin{aligned} cx_1^{m_1+m_2}x_{14}^{m_2} &= \lim_{x_i \rightarrow 0; 8, 10 \neq i \in \overline{2, 11}} x_1x_{14}(\partial_{x_1}\partial_{x_{14}} + \partial_{x_2}\partial_{x_{11}} + \partial_{x_3}\partial_{x_9} - \partial_{x_4}\partial_{x_7} + \partial_{x_5}\partial_{x_6})[x_1^{m_1} \\ &\quad \times (x_1x_{14} + x_2x_{11} + x_3x_9 - x_4x_7 + x_5x_6)^{m_2}] \\ &= m_2(m_1 + m_2 + 4)x_1^{m_1+m_2}x_{14}^{m_2} \end{aligned} \quad (3.99)$$

by (3.6)-(3.24), that is, $c = m_2(m_1 + m_2 + 4)$. We get:

Lemma 3.4. For $m_1, m_2 \in \mathbb{N}$,

$$\mathcal{D}_2(x_1^{m_1}\zeta_1^{m_2}) = m_2(m_1 + m_2 + 4)x_1^{m_1}\zeta_1^{m_2}. \quad (3.100)$$

According to Lemma 3.1,

$$V^4 = L(4, 0, 0) + L(2, 1, 0) + L(1, 0, 1). \quad (3.101)$$

Moreover, $L(1, 0, 1) = \eta V$. Thus the invariants in V^4W are $\mathbb{C}\eta\mathcal{D}_1$. Hence

$$[\mathcal{D}_2, \eta] = c_1\eta + c_2\eta\mathcal{D}_1 \quad \text{for some } c_1, c_2 \in \mathbb{C}. \quad (3.102)$$

Letting the above equation act on 1, we have

$$\mathcal{D}_2(\eta) = c_1\eta. \quad (3.103)$$

By (3.6)-(3.24) and (3.47),

$$\begin{aligned} 3c_1x_1x_{14}x_{27} &= \lim_{x_i \rightarrow 0; 14 \neq i \in \overline{2,16}} \eta = \lim_{x_i \rightarrow 0; 14 \neq i \in \overline{2,16}} \mathcal{D}_2(\eta) \\ &= 3(x_1x_{14}\partial_{x_1}x_{14} + x_{14}x_{27}\partial_{x_{14}}\partial_{x_{27}} + x_1x_{27}\partial_{x_1}x_{27})(x_1x_{14}x_{27}) = 9x_1x_{14}x_{27}, \end{aligned} \quad (3.104)$$

So $c_1 = 3$. Letting (3.102) act on x_1 , we have:

$$\mathcal{D}_2(\eta x_1) = (3 + c_2)\eta x_1. \quad (3.105)$$

As (3.104),

$$\begin{aligned} 3(3 + c_2)x_1^2x_{14}x_{27} &= \lim_{x_i \rightarrow 0; 14 \neq i \in \overline{2,16}} (3 + c_2)\eta x_1 = \lim_{x_i \rightarrow 0; 14 \neq i \in \overline{2,16}} \mathcal{D}_2(\eta x_1) \\ &= 3(x_1x_{14}\partial_{x_1}x_{14} + x_{14}x_{27}\partial_{x_{14}}\partial_{x_{27}} + x_1x_{27}\partial_{x_1}x_{27})(x_1^2x_{14}x_{27}) = 15x_1^2x_{14}x_{27}, \end{aligned} \quad (3.106)$$

Hence $c_2 = 2$. We get:

Lemma 3.5. *As operators on $\mathcal{A}$,*

$$[\mathcal{D}_2, \eta] = \eta(3 + 2\mathcal{D}_1). \quad (3.107)$$

For $m, m_1, m_2 \in \mathbb{N}$ with $m > 0$, we have

$$\begin{aligned} \mathcal{D}(\eta^m x_1^{m_1} \zeta_1^{m_2}) &= [m(111 + 11m_1 + m_2(m_1 + m_2 + 26)) \\ &\quad + \sum_{s=1}^m s(33 + 9(3s + m_1 + 2m_2))] \eta^{m-1} x_1^{m_1} \zeta_1^{m_2} \neq 0 \end{aligned} \quad (3.108)$$

by Lemmas 3.3-3.5. According to (3.84) and (3.108), we have:

Lemma 3.6. *For any $0 \neq f \in \mathcal{A}$,*

$$\mathcal{D}(\eta f) \neq 0. \quad (3.109)$$

The above lemma implies that

$$\{f \in \mathcal{A} \mid \mathcal{D}(f)\} = \sum_{m_1, m_2=0}^{\infty} L(m_1, m_2, 0). \quad (3.110)$$

Recall that $\mathcal{A}_m$ be the subspace of homogeneous polynomials of degree m in $\mathcal{A}$. Denote

$$\Phi_m = \{f \in \mathcal{A}_m \mid \mathcal{D}(f) = 0\}. \quad (3.111)$$

In summary, we have the following version of the main theorem.

Theorem 3.7. *The set $\{x_1^{m_1}\zeta_1^{m_2}\eta^{m_3} \mid n_1, m_2, m_3 \in \mathbb{N}\}$ is the set of all singular vectors in $\mathcal{A}$ up to a scalar multiple. In particular, η is the unique fundamental invariant (up to constant) and the identity*

$$(1 - q)^{26} \sum_{m_1, m_2=0}^{\infty} (\dim V(m_1\lambda_1 + m_2\lambda_6)) q^{m_1+2m_2} = 1 + q + q^2 \quad (3.112)$$

holds. Furthermore,

$$\mathcal{A}_k = \Phi_k \oplus \eta \mathcal{A}_{k-3} \quad \text{for } k \in \mathbb{N} \quad (3.113)$$

and

$$\Phi_m = \sum_{i=0}^{\lfloor m/2 \rfloor} L(m - 2i, i, 0) \quad \text{for } m \in \mathbb{N}, \quad (3.114)$$

where we treat $\mathcal{A}_r = \{0\}$ if $r < 0$.

References

- [1] J. Adams, *Lectures on Exceptional Lie Groups*, The University of Chicago Press Ltd., London, 1996.
- [2] G. Anderson and Blažek, E_6 unification model building.I. Clebsch-Gordan coefficients of $27 \otimes \overline{27}$, *J. Math. Phys.* **41** (2000), no. 7, 4808-4816.
- [3] G. Anderson and Blažek, E_6 unification model building.II. Clebsch-Gordan coefficients of $78 \otimes 78$, *J. Math. Phys.* **41** (2000), no. 12, 8170-8189.
- [4] G. Anderson and Blažek, E_6 unification model building.III. Clebsch-Gordan coefficients in E_6 tensor products of the 27 with higher-dimensional representations, , *J. Math. Phys.* **46** (2005), no. 1, 013506, 13pp.
- [5] M. Aschbacher, The 27-dimensional module for E_6 .I., *Invent. Math.* **89** (1987), no. 1, 159-195.
- [6] P. Berglund, P. Candelas, X. de le Ossa, E. Derrick, J. Distler and T. Hübsch, On instanton contributions to the masses and couplings of E_6 singlets, *Nuclear Phys. B* **454** (1995), no. 1-2, 127-163.
- [7] B. Binetruy and Zierau, A singular representation of E_6 , *Trans. Amer. Math. Soc.* **341** (1994), no. 2, 771-785.

- [8] J. Bion-Nadal, Subfactor of the hyperfinite II_1 factor with Coxeter graph E_6 as invariant, *J. Operator Theory* **28** (1992), 27-50.
- [9] R. Brylinski and B. Kostant, Minimal representations of E_6 , E_7 , and E_8 and the generalized Capelli identity, *Proc. Nar. Acad. Sci. U.S.A.* **91**, no. 7., 2469-2472.
- [10] B. Cao, Solutions of Navier equations and their representation structure, *arXiv: 0810.4766[math-ph]*.
- [11] B. Cerchiai and A. Scotti, Mapping the geometry of the E_6 group, *J. Math. Phys.* **49** (2008), no. 1, 012107, 19pp.
- [12] C. Das and L. Laperashvili, Preon model and family replicated E_6 unification, *SIGMA* **4** (2008), 012, 15pages.
- [13] L. Dickson, A class of groups in an arbitrary realm connected with the configuration of the 27 lines on a cubic surface, *J. Math.* **33** (1901), 145-123.
- [14] J. Fernández-Núñez, W. Garcia-Fuertes and A. Perelomov, Irreducible characters and Clebsch-Gordan series for the exceptional algebra E_6 : an approach through the quantum Calogero-Sutherland model, *J. Nonlinear Math. Phys.* **12**, suppl. 1, 280-301.
- [15] A. Ghezelbash, A. Shafiekhani and M. Abolbasani, On the Picard-Fuchs equations of $N = 2$ supersymmetric E_6 Yang-Mills theory, *Modern Phys. Lett. A* **13** (1998), no. 7, 527-531.
- [16] D. Ginzburg, On standard L -functions for E_6 and E_7 , *J. Reine Angew. Math.* **465** (1995), 101-131.
- [17] N. Haba and T. Matsuoka, Large lepton flavor mixing and E_6 -type unification models, *Progr. Theoret. Phys.* **99** (1998), no. 5, 831-842.
- [18] J. E. M. Homos and Y. M. M. Homos, Algebraic model for the evolution of the generic code, *Phys. Rev. Lett.* **71** (1991), 4401-4404.
- [19] R. Hubert, The (A_2, G_2) duality in E_6 , octonions and the triality principle, *Trans. Amer. Math. Soc.* **360** (2008), no. 1, 347-367.
- [20] R. Howl and S. King, Minimal E_6 supersymmetric standard model, *J. High Ener. Phys.* (2008) no. 1, 039, 31pp.

- [21] A. Iltyakov, On rational invariants of the group E_6 , *Proc. Math. Soc.* **124** (1996), no. 12, 3637-3640.
- [22] B. Kostant, On Macdonald's η -function formula, the Laplacian and generalized exponents, *Adv. Math.* **20** (1976), 179-212.
- [23] B. Kostant, Powers of the Euler product and commutative subalgebras of a complex simple Lie algebra, *Invent. Math.* **158** (2004), 181-226.
- [24] C. Luo, Noncanonical polynomial representations of classical Lie algebras, *arXiv: 0804.0305[math.RT]*.
- [25] I. Macdonald, Affine root systems and Dedekind's η -function, *Invent. Math.* **15** (1972), 91-143.
- [26] I. Morrison, P. Pieruschka and B. Wybourne, The interacting boson model with the exceptional groups G_2 and E_6 , *J. Math. Phys.* **32** (1991), no. 2, 356-372.
- [27] Y. Okamoto and R. Marshak, A grand unification preon model with E_6 metacolor, *Nuclear Phys. B* **268** (1986), no. 2, 397-405.
- [28] H. Rubenthaler, The (A_2, G_2) duality in E_6 , octonians and the triality principle, *Trans. Amer. Math. Soc.* **360** (2008), no. 1, 347-367.
- [29] K. Suzuki and M. Wakui, On the Turaev-Viro-Ocneanu invariant of 3-manifolds derived from the E_6 -subfactor, *Kyushu J. Math.* **56** (2002), 59-81.
- [30] J. Tits, A local approach to buildings, in *Geometric Vein*, Berlin-Heidelberg-New York, Springer, 1982, pp. 519-547.
- [31] X. Wang, Identification of Gepner's model with twisted LG model and E_6 singlets, *Modern Phys. Lett. A* **6** (1991), no. 23, 2155-2162.
- [32] X. Xu, Flag partial differential equations and representations of Lie algebras, *Acta Appl. Math.* **102** (2008), 149-280.